



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

of the roots of the given equation, and Lagrange's solution by continued fractions are also explained.

In Chap. XI determinants receive a clear natural treatment, while the subject of resultants and discriminants is carefully and rigorously discussed in the closing chapter.

On the whole in this book there is much to praise and little with which to find fault.

ELIJAH SWIFT.

## PROBLEMS AND SOLUTIONS.

B. F. FINKEL, CHAIRMAN OF THE COMMITTEE.

### PROBLEMS FOR SOLUTION.

**SPECIAL NOTICE.** In proposing problems and in preparing solutions, contributors will please follow the form established by the MONTHLY, as indicated on the following pages.

In particular, a solution should be preceded by the number of the problem, the name and address of the proposer, the statement of the problem, and the name and address of the solver.

The solution should then be given with careful attention to legibility, accuracy, brevity without obscurity, paragraphing and spacing, having in mind the form in which it will appear on the printed page.

Please use paper of letter size, write on one side only, leaving ample margins, put one solution only on a single sheet and include only such matter as is intended for publication.

Drawings must be made *clearly* and *accurately* and an extra copy furnished on a *separate sheet* ready for the engraver.

Unless these directions are observed by contributors, solutions must be entirely rewritten by the committee or else rejected.

Selections for this department are made two months in advance of publication.

Please send all solutions direct to the chairman of the committee.

MANAGING EDITOR.

### ALGEBRA.

When this issue was made up solutions of 410-19 had been received. Solutions of 406 and 420 are desired.

**420. Proposed by ELBERT H. CLARKE, Purdue University.**

Given the infinite series,

$$\frac{a}{r} + \frac{b}{r^2} + \frac{a+b}{r^3} + \frac{a+2b}{r^4} + \frac{2a+3b}{r^5} + \cdots,$$

in which  $a$  and  $b$  are any numbers and where each numerator after the first two is the sum of the two preceding numerators. To find the region of convergence and the sum of the series.

This problem is a generalization of one solved in the January number of the MONTHLY.

**421. Proposed by C. N. SCHMALL, New York City.**

Give a trigonometrical solution of the general quadratic equation.

### GEOMETRY.

When this issue was made up solutions of 437-38-39-40-43-45-47-48-54 had been received. Solutions of 427-30-32-33-44-46 are desired.

**449. Proposed by H. E. TREFETHEN, Colby College.**

Find a tetrahedron with the face angles at one vertex in arithmetical progression and its six edges expressed in integers.